

## **LITHOFLEX WORKSHOP**

**24-25 JUNE 2008**

**Research Centre Rotvoll –  
Trondheim – Norway**

### **Lithoflex theoretical background: Part IV: Isostatic anomalies and basin evolution**

**Jörg Ebbing  
Geological Survey of Norway  
NTNU Trondheim**

**with help of:  
Carla Braitenberg Christine Fichler, Laura  
Marello, Patrizia Mariani, Stephanie C.  
Werner, Susann Wienecke**



**StatoilHydro**



## **Isostatic anomalies and basin evolution**

- 1. Local isostasy**
  - Airy-Heiskanen
  - Pratt model
- 2. Regional isostasy (more in previous and next talk)**
- 3. Basin evolution (McKenzie-rifting, cratonic basins)**
- 4. Process oriented gravity modelling**
- 5. Backstripping**



**Isostasy** is a term used in geology to refer to the state of gravitational equilibrium between the Earth's lithosphere and asthenosphere such that the tectonic plates "float" at an elevation which depends on their thickness and density. It is invoked to explain how different topographic heights can exist at the Earth's surface. When a certain area of lithosphere reaches the state of isostasy, it is said to be in *isostatic equilibrium*.

<http://www.discoverourearth.org/student/topography/isostasy.html>



**Isostasy** is a term used in geology to refer to the state of gravitational equilibrium between the Earth's lithosphere and asthenosphere such that the tectonic plates "float" at an elevation which depends on their thickness and density. It is invoked to explain how different topographic heights can exist at the Earth's surface. When a certain area of lithosphere reaches the state of isostasy, it is said to be in *isostatic equilibrium*.

In general, the following types of isostatic concepts exist :

- Airy-Heiskanen isostasy
- Pratt isostasy
- Vening Meinesz isostasy

Isostatic anomalies represent in most cases Airy-Heiskanen compensation.




## Isostatic anomalies

$$g_{\text{Iso}} = g_{\text{BA}} - \Delta g_{\text{Iso}}$$

$$g_{\text{BA}} = g_{\text{observed}} - \Delta g_{\text{topo}} - \Delta g_{\text{BA}} - \gamma_0$$


$\Delta g_{\text{Iso}}$  = Isostatic reduction (Model)  
 $\gamma_0$  = Earth Normal gravity field  
 $\Delta g_{\text{topo}}$  = Topographic reduction  
 $\Delta g_{\text{BA}}$  = Bouguer plate reduction



## Concepts of local isostasy


### AIRY MODEL

VARIABLE DEPTH OF COMPENSATION  
 CONSTANT 'CRUSTAL' (LITHOSPHERIC) DENSITY




### PRATT MODEL

CONSTANT DEPTH OF COMPENSATION  
 VARIABLE 'CRUSTAL' (LITHOSPHERIC) DENSITY



TODAY, WE FEEL THAT THE ISOSTATIC MODEL IS A COMBINATION OF THE TWO THEORIES. EARTH'S RESPONSE TO LOADING DUE TO OROGENY, SEDIMENTATION, AND/OR GLACIATION INVOLVES A COMPLEX MASS-BALANCING INTERACTION BETWEEN THE RIGID LITHOSPHERE AND THE 'IMPRESSIONABLE' OR DEFORMABLE ASTHENOSPHERE.

## Continental lithosphere densities



Rock types:

BG = basaltic, gabbroic in oceanic crust

GAG = amphibolitic & granulitic in continental crust

PE = peridotitic, ultramafic

D = Dioritic (?), possibly amphibolitic

S = sediments

GR = granitic gneissic upper crust


W = water

P-wave velocities (km/s):  
black numbers



Densities (g/cm³):  
White numbers

## Airy isostasy

- Collisional orogens
- Rift zones
- Non-volcanic margins (e.g. Iberia)





## Eastern Alps: Topography and Bouguer anomaly



Ebbing 2002




## Eastern Alps: Topography and Moho



Ebbing 2002




## TRANSALP



Ebbing et al. 2006


## Pratt isostasy

- Fennoscandia
- Volcanic continental margins  
– mid-Norway






## Topography and gravity




Topography/bathymetry compilation:  
Dehls et al. 2000



Gravity map as compiled by:  
Skilbrei et al. (2000), Korhonen et al. (2002)




## Seismic and gravity studies



### Deep seismic reflection and refraction experiments


Seismic Moho depth after Kinck et al. 1993,  
Mjelde et al. 2005, Olesen et al. 2002




Gravity map as compiled by:  
Skilbrei et al. (2000), Korhonen et al. (2002)



## Pratt isostasy and gravity residuals



Isostatic high-density lower crust




Gravity residual of isostatic model

Ebbing, 2007





## Mid-Norwegian margin




Ebbing et al. 2008








## Isostatic state of the margin




Seismic Moho




Isostatic Moho



## Origin and extent of LCB?



Ebbing et al. , in press



Mjelde et al. 2008


New interpretation: magmatic  
underplating (yellow) + eclogite (red)



## Isostasy and flexural rigidity

### Local Isostasy - model after Airy

$$t_i(\mathbf{x}, \mathbf{y}) = -T_0 - \frac{\rho_c}{\rho_m - \rho_c} h_i(\mathbf{x}, \mathbf{y})$$



### Regional Isostasy - model after Vening Meinesz



## Isostasy and flexural rigidity Vening Meinesz

Flexural models or models of a thin elastic plate use the transfer function

$$W(\mathbf{k}) = -\frac{\rho_c}{\rho_m - \rho_c} \left( 1 + \frac{16\pi^4 \mathbf{k}^4 D}{(\rho_m - \rho_c)g} \right)^{-1} H(\mathbf{k})$$


and

$$t_i^*(\mathbf{x}, \mathbf{y}) = -T_0 - w_i(\mathbf{x}, \mathbf{y})$$


$D$  = flexural rigidity of the lithospheric plate

$H(\mathbf{k}), W(\mathbf{k})$  = Fourier Transformation of topography and deflection

$w(\mathbf{x}, \mathbf{y})$  = Inverse Fourier Transformation of  $W(\mathbf{k})$



## Gravity field after regional isostatic correction





Olesen et al. 2007




## Isostasy and basin modelling

- Simple rift basin  
McKenzie type ~Airy-Heiskanen isostasy
- Extensional margins  
Vøring-Viking
- Cratonic and intra-cratonic basins  
Barents Sea






Isostatic response to loading and its gravity signal is depending on:  
1) density (contrast)



Karner et al., 2005




Isostatic response to loading and its gravity signal is depending on:  
1) density (contrast) and 2) strength of the lithosphere





Isostatic response to loading and its gravity signal is depending on:  
1) density (contrast) and 2) strength of the lithosphere



time-constant  $T_e$ :  
water loading

time-constant  $T_e$ :  
sediment loading


time-variable  $T_e$ :  
Sediment loading

Karner et al., 2005



Isostatic response to loading and its gravity signal is depending on:  
1) density (contrast) and 2) strength of the lithosphere. We can use this for:


### Process oriented gravity modelling



Karner et al., 2005




Crustal extension and isostatic response is connected to stretching factors




Karner et al., 2005

The  $T_e$  distribution we apply to create the present-day gravity signal can be used and tested in backstripping




Kusznir et al. 2004

## Basin modelling: backstripping




Roberts et al. 1997

## Typical rift basin




## Non-typical rift basin, e.g. cratonic basin



...or after thermal post-rift subsidence




## The crustal structure of the Barents Sea




Bungum et al. 2005; Ritzmann et al. 2007



### 3D density modelling: model set up from Barents3D




### 3D density modelling Gravity effect of Barents50 - Profil




- Possible error sources:**
- Geometry of Barents50
  - Densities of Barents50

## Isostatic considerations Isostatic vs. seismic Moho



**Airy Moho  
(incl. Sediment loading)**



**Seismic Moho**


Ebbing et al. 2007





## Isostatic density inversion

- Gravity
  - Regional trend/offset
- Isostasy
  - Moho is not compensating for loading by sediments & topography

=> Isostatic density inversion







## Summary

- isostatic concepts allow to model lithospheric structures
- isostatic modelling allows feedback on internal mass distribution and evolution
- concept oriented gravity modelling allows to evaluate different scenarios for basin development
- A careful selection of the isostatic concept (Airy-Heiskanen, Pratt, Vening Meinesz) has always to be made
- Flexural isostasy can also give information on crustal rheology (*not covered here*)



## Literature 1/2

- Allen, P.A. and Allen, J.R., 2005. *Basin Analysis: Principles and Applications*, 2. ed., Blackwell Publishing, 549 pp.
- Birch F. (1991). The velocity of compressional waves in rocks to 10 kbar. Part 2', *J. Geophys. Res.* 66, 2199-2224.
- Braitenberg, C., Wienecke, S. and Wang, Y., 2006. Basement structures from Satellite Derived Gravity Field: the South China Sea Ridge. *Journal of Geophysical Research-Solid Earth*, 111, B05407.
- Brocher, T. M. 2005 Empirical Relations between Elastic Wavespeeds and Density in the Earth's Crust *Bulletin of the Seismological Society of America*; December 2005; v. 95; no. 6; p. 2081-2092
- Bungum, H., Ritzmann, O., Maercklin, N., Faleide, J., Mooney, W.D. & Detweiler, S.T. 2005: Three-Dimensional Model for the Crust and Upper Mantle in the Barents Sea Region. *EOS*, 86(16), doi: 10.1029/2005EO160003.
- Carlson R.L. and Herrick C.N. (1990). Densities and porosities in the oceanic crust and their variations with depth and age, *J. Geophys. Res.*, 95, 9153-9170.
- Carmichael, R.S. 1984. *Handbook of physical properties of rocks*. Volume I (1986), II (1982), III (1984).
- Christensen, N. I., and W. D. Mooney (1995). Seismic velocity structure and composition of the continental crust: A global view, *J. Geophys. Res.* 100, 9761-9788.
- Dehls, J.F., Olesen, O., Bungum, H., Hicks, E., Lindholm, C.D. & Riis, F. 2000: Neotectonic map, Norway and adjacent areas 1:3 mill. Norges geologiske undersøkelse, Trondheim, Norway.
- Ebbing, J., 2002. 3D-Dichtemodellierung und isostatisches Verhalten der Lithosphäre in den Ostalpen (The 3D density structure and isostatic compensation of the lithosphere in the Eastern Alps). German *Ph.D thesis*, Freie Universität Berlin.
- Ebbing, J., 2007: Isostatic density modelling explains the missing root of the Scandes. *Norwegian Journal of Geology*, 87, 13-20.
- Ebbing, J., Braitenberg, C. and Götze, H.-J., 2006. The lithospheric density structure of the Eastern Alps. *Tectonophysics*, 414, 145-155.
- Ebbing, J., Braitenberg, C. and Wienecke, S., 2007. Insights into the lithospheric structure and the tectonic setting of the Barents Sea region from isostatic considerations. *Geophysical Journal International*, 171 (3), 1390-1403.
- Ebbing, J., Skilbrei, J.R. and Olesen, O. 2007. Insights into the magmatic architecture of the Oslo Graben by petrophysically constrained analysis of the gravity and magnetic field, *J. Geophys. Res.*, 112, B04404, doi:10.1029/2006JB004694.
- Ebbing J., Olesen, O., Gernigon, L., Reynisson, R.F., og sokkelgeofysikk lag på NGU 2008: Tyngde og magnetiske data viser sporene av gamle strukturer på norsk sokkel. I: *Geologi for samfunnet i 150 år - arven etter Kjerulf* (T. Slagstad & R. Dahl, eds.), *Gråsteinen* 12, 89-98.
- Ebbing, J., Gernigon, L., Pascal, C., Olesen, O. and Osmundsen, P.T., A discussion of structural and thermal control of magnetic anomalies on the mid-Norwegian margin. *Geophysical Prospecting*, in press.
- Faleide, J.I., Ritzmann, O., C. Weidle, C. & A. Levshin, A. 2006: Geodynamical aspects of a new 3D geophysical model of the greater Barents Sea region – Linking sedimentary basins to the upper mantle structure. *Geophysical Research Abstracts*, 8, 08640.
- Gardner, G. H. F., L. W. Gardner, and A. R. Gregory (1984). Formation velocity and density -The diagnostic basics for stratigraphic traps, *Geophysics* 39, 770-780.
- Godfrey, N. J., B. C. Beaudoin, S. L. Klemperer, and the Mendocino Working Group USA (1997). Ophiolitic basement to the Great Valley forearc basin, California, from seismic and gravity data: Implications for crustal growth at the North American continental margin, *Geol. Soc. Am. Bull.* 109, 1536-1562.
- Karner, G.D., Studinger, M., Bell, R.E., 2005. Gravity anomalies of sedimentary basins and their mechanical implications: Application to the Ross Sea West Antarctica. *Earth and Planetary Science Letters*, 235( 3-4), 577-596.



## Literature 2/2

- Kinck, J.J., Husebye, E.S. & Larsson, F.R. 1993: The Moho depth distribution in Fennoscandia and the regional tectonic evolution from Archean to Permian times. *Precambrian Research* 64, 23-51.
- Korhonen, J., V., Aaro, S., All, T., Elo, S., Haller, L.Ä., Kääriäinen, J., Kulnich, A., Skilbrei, J.R., Solheim, D., Säävuori, H., Vaher, R., Zhdanova, L. & Koistinen, T. 2002a. Bouguer anomaly map of the Fennoscandian shield 1: 2 000 000. Geological Surveys of Finland, Norway and Sweden and Ministry of Natural Resources of Russian Federation.
- Kusznir, N.J., Hunsdale, R. and Roberts, A.M., 2004. Timing of depth-dependent lithosphere stretching on the S.Lofoten riftedmargin offshoremid-Norway: pre-breakup or post-breakup?, *Basin Research* 16, 279–296.
- Larsen, H.C., Saunders A.D., Clift P.C., and the Shipboard Scientific Party (1994). Proceeding of the Drilling Program Initial Report. Vol. 152, Ocean Drill. Program. College Station, Tex.
- Ludwig W.J., Nafe J.E. and Drake C.L. (1970). Seismic refraction, in the Sea, A.E.Maxwell (Editor), Vol.4, Wiley-Interscience, New York, 52-84
- Meissner, R., 1986: The continental crust: A geophysical approach. Academic Press Inc., p. 4
- Mjelde, R., Raum, T., Breivik, A., Shimamura, H., Murai, Y., Takanami, T. & Faleide, J.I. 2005: Crustal structure of the Vøring margin, NE Atlantic: a review of geological implications based on recent OBS-data. In: Doré, A.G. & Vining, B.A. (eds.) *Petroleum Geology: North-West Europe and Global Perspectives*, Proceedings of the 6th Petroleum Geology Conference. Geological Society Special Publication, Geological Society, London, 803-813.
- Mjelde, R., Faleide, J.I., Breivik, A.J., Raum, T., Lower crustal composition and crustal lineaments on the Vøring Margin, NE Atlantic: A review, *Tectonophysics* (2008), doi: 10.1016/j.tecto.2008.04.018.
- Mussett, A. E., Khan, M.A., 2000. Looking into the earth. An introduction to Geological Geophysics. Cambridge University Press
- Olesen, O., Lundin, E., Nordgulen, Ø., Osmundsen, P.T., Skilbrei, J.R., Smethurst, M.A., Solli, A., Bugge, T. & Fichler, C. 2002: Bridging the gap between the onshore and offshore geology in Nordland, northern Norway. *Norwegian Journal of Geology* 82, 243-262.
- Olesen, O., Dehls, J.F., Ebbing, J., Kihle, O. & Lundin, E., 2007. Aeromagnetic mapping of deep-weathered fracture zones in the Oslo Region – a new tool for improved planning of tunnels. *Norwegian Journal of Geology*, 87, 253-267.
- Ritzmann, O., Maercklin, N., Faleide, J.I., Bungum, Mooney, W.D. & Detweiler, S.T. 2007: A three-dimensional geophysical model of the crust in the Barents Sea region: model construction and basement characterization. *Geophys. J. Int.* 170, 417–435.
- Roberts, A.M., Lundin, E. and Kusznir, N.J. 1997. Subsidence of the Vøring Basin and the influence of the Atlantic continental margin. *Journal of the Geological Society, London*, 154, 1997, 551–557.
- Skilbrei, J.R., Kihle, O., Olesen, O., Gellein, J., Sindre, A., Solheim, D. & Nyland, B., 2000: Gravity anomaly map Norway and adjacent ocean areas, scale 1:3 Million, Geological Survey of Norway, Trondheim.
- Sobolev, S. and Babeyko, A., 1994. Modeling of mineralogical composition, density and elastic wave velocities in anhydrous magmatic rocks. *Surveys in Geophysics*, 15: 515-544.
- Tsikalas, F., 1992: *A study of seismic velocity, density and porosity in Barents Sea wells (N. Norway)*. Master Thesis, Department of Geology, University of Oslo, Norway.
- Wienecke, S., 2006. A new analytical solution for the calculation of flexural rigidity: significance and applications. PhD Thesis, Free University Berlin, Germany. <http://www.diss.fu-berlin.de/2006/42/indexe.html>.
- Wienecke, S., Ebbing, J. & Gernigon, L. 2007: 3D density modelling, isostasy and elastic thickness calculation of the Barents Sea. NGU Report 2007.022. 56pp.

